



UW PACC

Psychiatry and Addictions Case Conference

UW Medicine | Psychiatry and Behavioral Sciences

LONELINESS – WHY IT IS IMPORTANT, HOW TO ASK ABOUT IT AND HOW TO HELP

**SEBASTIAN TONG, MD, MPH
DEPARTMENT OF FAMILY MEDICINE,
UNIVERSITY OF WASHINGTON
APRIL 23, 2026**



SPEAKER DISCLOSURES

- ✓ Funding from National Institutes of Health (R34DA058325). Opinions here are my own and do not reflect the perspectives of NIH or HHS.
- ✓ No other relevant disclosures

PLANNER DISCLOSURES

The following series planners have no relevant conflicts of interest to disclose; other disclosures have been mitigated.

Mark Duncan MD

Rick Ries MD

Kari Stephens PhD

Barb McCann PhD

Anna Ratzliff MD PhD

Betsy Payn MA PMP

Esther Solano

Cara Towle MSN RN

OBJECTIVES

1. To describe the health risks and comorbidities associated with loneliness.
2. To identify tools to screen for loneliness in health care settings.
3. To identify ways to address loneliness in health care settings.

PATIENT #1

68-year-old Amharic speaking female with hypertension who presents for routine follow-up complaining of worsening chronic wrist, shoulder and knee pain. She was referred 2 months ago to physical therapy but did not go.

PATIENT #2

23-year-old male presenting to establish care asking for ADHD medications to help with concentrating at work.

WHY SHOULD WE CARE?

Our Epidemic of Loneliness and Isolation



2023

The U.S. Surgeon General's Advisory on the Healing Effects of Social Connection and Community



A CRITIC AT LARGE

A.I. IS ABOUT TO SOLVE LONELINESS. THAT'S A PROBLEM

The discomfort of loneliness shapes us in ways we don't recognize—and we may not like what we become without it.

By Paul Bloom

July 14, 2025

New York Times Magazine | Why Is the Loneliness Epidemic So Hard to Cure?

MENTAL HEALTH

DEFINITIONS

Loneliness

- Subjective
- Distressing feeling from a perceived gap or deficit in social relationships
- Not the same thing as being shy or introverted

Social isolation

- Objective
- State of being physically separate from others
- Lack of social contact or relationships

DEFINITIONS

LONELINESS



Feelings of distress related to lack of desired connections

SOCIAL ISOLATION




Physically alone but not necessarily distressed

PREVALENCE

- **Loneliness:** More than ½ U.S. adults
 - Young adults ≤ 35 years old may have highest rates of loneliness
 - Older adults have many known risk factors
- **Social Isolation:** 24% of community-dwelling older adults

FRAMEWORK



NEGATIVE HEALTH IMPACTS


- Physical
- Cognition
- Mental
- Lifespan



HEALTH IMPACTS: PHYSICAL

Loneliness and social isolation
increase the risk of:

- Stroke and cardiovascular disease
- Hypertension
- Type 2 Diabetes
- Poor sleep quality
- Physical inactivity and worsened function
- and more!



HEALTH IMPACTS: BRAIN/COGNITION

- Loneliness and social isolation increase the risk of:
 - Faster cognitive decline
 - Overall worse memory and executive function
 - Mild Cognitive Impairment (MCI)
 - Dementia, including Alzheimer's Disease

50%

Chronic social isolation and loneliness can increase the risk of developing dementia by **50%** in older adults

HEALTH IMPACTS: MENTAL HEALTH

- Loneliness and social isolation increase the risk of:
 - Depression and anxiety
 - Suicidal thoughts
 - Decreased quality of life
 - Deaths of despair (alcohol, drug overdose, suicide)




HEALTH IMPACTS: LIFESPAN

- Loneliness increases risk for early death by 26%
- Social isolation increases risk for early death by 29%



The Three Vital Components of Social Connection


The extent to which an individual is socially connected depends on multiple factors, including:



ELEMENTS OF LONELINESS

Element of Loneliness	Description	Examples	Possible Consequences
Emotional Loneliness	Absence of close emotional attachment or intimate relationship. Often linked to loss (e.g., breakup, death, separation).	Missing a confidant, partner, or close friend.	Depression, grief, feelings of emptiness.
Social Loneliness	Lack of a broader social network or sense of belonging to a group.	Few friends, weak community ties, isolation at school or work.	Social withdrawal, anxiety, low self-esteem.
Existential Loneliness	A deeper sense of isolation stemming from awareness of life's meaninglessness, mortality, or disconnection from humanity.	Feeling fundamentally separate even when surrounded by others.	Despair, spiritual questioning, existential anxiety.

NASEM FRAMEWORK FOR LONELINESS



HOW DO WE IDENTIFY LONELINESS?

LONELINESS

- UCLA Loneliness Scale (20 or 3-item version)
 - How often someone lacks companionship, feels left out, or feels isolation
- De Jong Gierveld Scale (11 or 6-item version)
 - Includes emotional and social loneliness subscales
- Single-item measure
 - How often someone feels lonely (often/always, some of the time, occasionally, hardly ever, never)
 - Concern about potential stigma; meta-analyses illustrates lower prevalence of chronic loneliness

SOCIAL ISOLATION

- 50+ measures, no consensus
- Lubben Social Network Scale (LSNS, 6-item version)
 - Captures social networks, social isolation, social support
 - Derived from the BSNI to be more specific to older adults
- Berkman-Syme Social Network Index (BSNI, 4-item version)
 - Measures frequency and quality of social support, social participation, and social contact
 - Recommended for inclusion in EHR by prior IOM committee

SCREENING IN PRIMARY CARE


Use screening tools to identify loneliness and social isolation

- Universal screening at predetermined intervals (e.g. annual exams)
- Targeted screening for high-risk individuals (e.g. bereavement, mobility or sensory impairments, transportation insecurity, major life transitions)

If positive, provide education on the negative impacts of loneliness and social isolation

- Perform in-depth assessment and address any contributors
- Leverage other team members (case managers, social workers)
- Offer tailored referrals

POTENTIAL SCREENING PRACTICE IN CLINICAL SETTINGS



WHAT CAN WE DO IN PRACTICE TO REDUCE LONELINESS?

FRAMEWORK FOR INTERVENTIONS

The Six Pillars to Advance Social Connection



1

Strengthen Social Infrastructure in Local Communities

Design the built environment to promote social connection

Establish and scale community connection programs

Invest in local institutions that bring people together

2

Enact Pro-Connection Public Policies

Adopt a "Connection-in-All-Policies" approach

Advance policies that minimize harm from disconnection

Establish cross-departmental leadership at all levels of government

3

Mobilize the Health Sector

Train health care providers

Assess and support patients

Expand public health surveillance and interventions

4

Reform Digital Environments

Require data transparency

Establish and implement safety standards

Support development of pro-connection technologies

5

Deepen Our Knowledge

Develop and coordinate a national research agenda

Accelerate research funding

Increase public awareness

6

Build a Culture of Connection

Cultivate values of kindness, respect, service, and commitment to one another


Model connection values in positions of leadership and influence

Expand conversation on social connection in schools, workplaces, and communities

EVIDENCE

- Masi et al (2011) meta-analyses:
 - Improving social skills
 - Enhancing social support
 - Increasing opportunities for social contact
 - Addressing maladaptive social cognition

GROUP INTERVENTIONS



Social Prescribing:

- Aims to map out social networks and create goals for social connection
- Modeled from UK health service of “social prescribing”
- Groups include open discussion, problem solving, and resource sharing



Cognitive Behavioral Therapy:

- Adapted from a program created by Swedish researchers
- Focus on behavioral activation and managing thoughts and feelings
- More structured modules with practice examples

OUR ADAPTED SOCIAL PRESCRIBING INTERVENTION

- 8-week virtual, group intervention
- Activities:
 - Mapping my world (People, Places, Activities)
 - Connections Plan (goals, concrete steps, timeframe)
 - Spotlight activities
 - Social connection framework
 - Social anxiety
 - Letter to self




OUR ADAPTED CBT INTERVENTION

- 6-week virtual, group intervention
- Based on psychological, cognitive-behavioral therapy (CBT) techniques
- Address behaviors (e.g., avoidance) and cognitions associated with loneliness
- Incorporate with examples and techniques that are specific to chronic pain experience and management

Week 1	Engagement and Introduction to the CBT model of loneliness
Week 2	Identify your behaviors that are linked to loneliness
Week 3	Identify values, goals and planned behaviors that help improve loneliness
Week 4	Problem-solve what gets in the way of you connecting with others
Week 5	Working with your thoughts
Week 6	How to maintain improvements and new behaviors


RESULTS: LONELINESS SCORE

Change in UCLA-20 Scores Over Time




RESULTS: DEPRESSION

Change in PHQ-9 Scores Over Time



RESULTS: ANXIETY

Change in GAD-7 Scores Over Time



LONELINESS AND SOCIAL ISOLATION INTERVENTIONS

- Consider three overarching strategies:
 - #1) Promote social contact
 - #2) Transfer knowledge and skills to engage in social activity
 - #3) Address cognitive or psychological barriers for social contact (e.g., cognitive behavioral therapy)
- Use a patient-centered approach
- Involve other clinical care team members (case managers, social workers, behavioral health)
 - Consider interventions within healthcare system (therapy) and outside (referrals to community programs)
- Address barriers to engagement (chronic comorbidities, mobility or sensory impairment, etc.)

BACK TO OUR PATIENTS

- Case #1: 68-year-old female who is a full-time caregiver
- Case #2: 23-year-old male with concern for ADHD



UW PACC

Psychiatry and Addictions Case Conference

UW Medicine | Psychiatry and Behavioral Sciences

QUESTIONS?

SEBASTIAN TONG, MD, MPH

SETONG@UW.EDU

