



UW PACC

Psychiatry and Addictions Case Conference

UW Medicine | Psychiatry and Behavioral Sciences

EATING DISORDERS

MEGAN RIDDLE, MS MD PHD
(SHE/HER)



SPEAKER DISCLOSURES

- ✓ I work at the Eating Recovery Center in Bellevue
 - We take Medicaid (WA & OR)
 - I'm always happy to talk about eating disorders!
Please reach out
 - Megan.Riddle@ERCPathlight.com
 - Riddlem2@uw.edu
 - Office: 425-214-9335

OBJECTIVES

1. Discuss why eating disorders matter
2. Learn how to screen for and diagnose eating disorders and understand why this is important
3. Outline of treatment, including where to treat and when to refer out
4. Explore special considerations of BED

WHY TALK ABOUT EATING DISORDERS

- Eating disorders have a high morbidity and mortality
 - All eating disorders carry increased risk of death, both as a result of suicide and medical complications
- With 30 million people in the US with eating disorders, you will see these patients regardless of where you practice or what your focus is

WHY TALK ABOUT EATING DISORDERS?


- These patients fall through the cracks
 - We hold certain assumptions about who has eating disorders

WHY TALK ABOUT EATING DISORDERS?


- Medical education perpetuates our cultural beliefs about food, weight, shape and health
 - We may worsen an underlying eating disorder by recommending weight loss strategies to someone struggling with disordered eating – these can lead to worsening of eating behaviors and/or avoidance of the medical system

EATING DISORDERS

Anorexia nervosa

- Restriction of energy intake leading to low body weight
 - Intense fear of weight gain or behaviors that interfere with weight gain
 - Excess concerns about weight and shape
- 
- A faint, light blue illustration of a person standing on a scale, positioned behind the bulleted text for Anorexia nervosa.

Bulimia nervosa

- Recurrent binge episodes associated with eating an excess amount in a discrete period and lack of control
 - Compensatory behaviors to prevent weight gain
 - Binge eating and purging occur on average at least weekly for 3mo
 - Excess concerns about shape and weight
- 
- A faint, light blue illustration of a person sitting at a table and eating, positioned behind the bulleted text for Bulimia nervosa.

EATING DISORDERS

Binge Eating Disorder

- Recurrent binge episodes associated with eating an excess amount in a discrete period and lack of control
- **No** compensatory behaviors to prevent weight gain
- Binge eating occurs on average at least weekly for 3mo

Avoidant/Restrictive Food Intake Disorder

- Eating disturbance (i.e. concern about eating certain food types/textures) that prevents patient from meeting nutritional needs leading to:
 - Weight loss
 - Nutrient deficiency
 - Dependence on supplements/feeding tube
 - Interference with psychosocial
- Not in the setting of AN or BN or explained by other medical condition

EATING DISORDERS

Other Specified Feeding & Eating Disorder

- Symptoms of an eating disorder that cause clinically significant distress or impairment but do not meet the full criteria for any of the other disorders
- i.e. Atypical anorexia, purging disorder, BN or BED of short duration/ low intensity

Unspecified Eating Disorder

- Symptoms of an eating disorder that cause clinically significant distress are present but diagnosis is unclear

WHO TO SCREEN?

- American Academy of Pediatrics advocates the routine use of screening questions for all preteen and adolescent patients

WHO TO SCREEN?


- In adults, there aren't specific guidelines
- Screen high risk groups
 - Young adults
 - Those in the LGBTQ community, particularly genderqueer individuals
 - Female-identified and genderqueer individuals under stress, with anxiety
 - Everyone with a family history of eating disorders
 - Everyone with rapid changes in weight or those seeking help with weight loss
 - Athletes

EATING DISORDERS: QUICK SCREEN


- Screen for Disordered Eating
 - Do you often feel the desire to eat when you are emotionally upset or stressed?
 - Do you often feel that you can't control what or how much you eat?
 - Do you sometimes make yourself throw up (vomit) to control your weight?
 - Are you often preoccupied with a desire to be thinner?
 - Do you believe yourself to be fat when others say you are thin?
- Two positive responses gives sensitivity 90.5% and specificity 57.5%

(Maguen et al, 2018)


MAKING A DIAGNOSIS



MAKING A DIAGNOSIS




TREATMENT TEAM



WHERE TO TREAT

- Whether a patient should be hospitalized for treatment depends on several factors
 - Medical stability
 - Comorbid psychiatric issues
 - Willingness to engage in treatment



There are no specialized ED inpatient units in WA

MEDICAL EVALUATION

- These patients can have serious medical complications secondary to starvation or binge/purge behaviors
- Don't make assumptions about medical stability based on weight

MEDICAL ASSESSMENT

- Thorough physical exam
- Labs:
 - CBC, BMP, Mg, Phos, LFTs, TSH, free T4, Utox, UPreg
 - Consider: pre-albumin, urine electrolytes (diuretic use), amylase
- EKG
- Long-term weight-suppressed: DEXA scan


FACTORS SUPPORTING HOSPITALIZATION

	Adults	Adolescents (12-19 years)
Heart rate	<50 bpm	<50 bpm
Orthostatic change in heart rate	Sustained increase of >30 bpm	Sustained increase of >40 bpm
Blood pressure	<90/60 mmHg	<90/45 mmHg
Orthostatic blood pressure	>20 mmHg drop in sBP	>20 mmHg drop in sBP
Glucose	<60 mg/dl	
Potassium	Hypokalemia	Hypokalemia
Sodium	Hyponatremia	Hyponatremia
Phosphate	Hypophosphatemia	Hypophosphatemia
Magnesium	Hypomagnesemia	Hypomagnesemia
Temperature	<96.0 F	<96.0 F
BMI	<15	<75% of median BMI for age and sex
Rapidity of weight change	Greater than 10% decrease in body weight within the last 30 days	Greater than 10% weight loss in 6 months or greater than 20% weight loss in 1 year
Compensatory behaviors	Occur multiple times daily and have either caused severe physiological consequences	Occur multiple times daily and have either caused severe physiological

FACTORS SUPPORTING STEP-UP

- When to consider referral to a specialized center?
 - Medical instability – may need to be medically hospitalized first
 - Inability or unwillingness to change behaviors independently
 - Outpatient care not leading to a decrease in symptom frequency
 - Lack of access to specialized providers in the outpatient setting

TREATMENT




ADDRESS MALNOURISHMENT & ED BEHAVIORS

- Regardless of body shape and size, the majority of ED patients need nutritional rehab
 - Early in treatment, we work on identifying a target weight – based on weight history and trends
 - Target weight may or may not be near someone's “ideal” body weight
- Work with an RD to create a meal plan

ENCOURAGE THERAPY

- AN: Nutritional counseling + Therapy is better than nutritional counseling alone – no one type of therapy is superior
- BN & BED: Good evidence that Cognitive Behavioral Therapy is effective intervention
- Therapy for psychiatric comorbidities, ie. trauma



CONSIDER MEDICATIONS

- Anorexia nervosa
 - Medications have limited efficacy at low weight
 - No FDA approved meds
 - Antidepressants may help prevent relapse
 - Antipsychotics have mixed evidence
- Bulimia nervosa
 - SSRIs are 1st line: Fluoxetine is FDA approved
 - Avoid bupropion due to increased seizure risk
- Binge eating disorder
 - SSRIs are 1st line
 - Lisdexamfetamine (Vyvanse) is FDA approved*

RECOVERY

- Recovery has its ups and down

CAREGIVER BURDEN

- Caring for these patients is incredibly emotionally taxing for loved ones and clinicians
- These cases can bring up very intense emotions
- Creating space to acknowledge this is an important part of this work



UW PACC

Psychiatry and Addictions Case Conference

UW Medicine | Psychiatry and Behavioral Sciences

WHEN IT'S NEVER ENOUGH

CARING FOR INDIVIDUALS WITH BINGE EATING DISORDER



Huge thanks to Lindsay Birchfield, MS, RD, CD

EPIDEMIOLOGY



3.5%

of American Women
have BED



2%

of American Men
have BED




1.6%

of Adolescents
have BED

- Mean age of onset 23 - 25yo (older than BN or AN)
- Equally distributed across racial and ethnic groups





EPIDEMIOLOGY

- Not all obese patients have BED and not all BED patients are obese
- 36-42% of individuals with BED are obese
- Up to a quarter of patients with obesity struggle with BED

(Montano et al. 2016)


ETIOLOGY OF BINGE EATING DISORDER



ETIOLOGY OF BINGE EATING DISORDER


<http://binow.ru/wp-content/uploads/2013/09/razmyshleniya-o.jpg>

ETIOLOGY OF BINGE EATING DISORDER



<http://binow.ru/wp-content/uploads/2013/09/razmyshleniya-o.jpg>

ETIOLOGY OF BINGE EATING DISORDER



<http://binow.ru/wp-content/uploads/2013/09/razmyshleniya-o.jpg>

ETIOLOGY OF BINGE EATING DISORDER



<http://binow.ru/wp-content/uploads/2013/09/razmyshleniya-o.jpg>

ETIOLOGY OF BINGE EATING DISORDER




<http://binow.ru/wp-content/uploads/2013/09/razmyshleniya-o.jpg>


ADDRESSING BED

- BED treatment – maybe more so than other ED – really requires a multipronged approach that also openly addresses challenging topics like social inequities and personal/institutional/societal trauma in addition to beliefs about food and body shape
 - Honestly – we probably need to do more this in all our ED work (and all of mental health?) but it's a slow evolution

SATTER'S HIERARCHY OF FOOD NEEDS




CYCLE OF BINGE EATING



“In short, food becomes one of the only successful ways to simultaneously numb and soothe and enact self-loathing. It’s the perfect trifecta for someone to cope with a body that is deemed incorrect by the medical field and media.”
– Lindsay Birchfield, MS, RD, CD


INTERRUPTING THE CYCLE



INTERRUPTING THE CYCLE: CONSISTENT AND ATTUNED REGULAR EATING (CARE)

- Establish a pattern of Regular Eating
 - Emphasis on consistent eating at rhythmic intervals irrespective of appetite cues, motivation, and mood
 - Mechanical eating
 - Eating by the clock
 - Benefits include physical safety and stability, reduces deprivation –sensitive binges, reducing obsessive thoughts about food, helps restore hunger/satiety cues etc.
- 3 meals and 2-3 snacks/day, preplanned at designated times

INTERRUPTING THE CYCLE



UNMET NEEDS & EMOTIONAL TRIGGERS

- Work on identifying unmet needs
- Increase skill use
- Build social connections – including fat-affirming spaces
- Gentle nutrition and joyful movement to support wellbeing
- Mindful eating to promote emotional satisfaction
- Creating space, apart from food, for patient to experience feelings

GLP1AS AND EATING DISORDERS

- Lots of questions in the field regarding the interplay between EDs and GLP1as
- GLP1as can exacerbate ED behaviors or be misused as part of an ED
 - Patients with active EDs getting GLP1as for weight loss
 - From their provider who is unaware of their active ED, from online clinics (may misrepresent medical history), or off the black market
 - For patients with an ED history, GLP1as can lead to a relapse due mimicking ED behaviors (restriction, weight loss, sense of control)

GLP1AS AND EATING DISORDERS

- Could GLP1as have a role in ED treatment for binge eating?
 - Maybe? Research is still needed
 - Risk at not healing the relationship with food or addressing the underlying drivers of binge behavior, which may shift to something else

GLP1AS AND EATING DISORDERS: BEST PRACTICES

- Screen all patients who are being started on GLP1as for EDs
- Ask all individuals with EDs about use of GLP1as
- What about for patients with a history of EDs who want to start a GLP1a for medical comorbidities?
 - Develop a multidisciplinary team (Physician, RD, therapist)
 - Create contingency/exit plan: ie criteria for discontinuation
 - Carefully discuss risks/benefits, including risk of ED relapse

CONCLUSIONS

- BED is the most common ED and stems from a variety of biological, psychological and social factors – it's not about will power
- Consider where someone is in their own food hierarchy
- Treatment focuses less on the binge behavior and more on the contributing factors

THANK YOU!

- Thoughts or questions?